

Non-universal corrections to the level curvature distribution beyond random matrix theory.

I. V. Yurkevich^{1,3} and V. E. Kravtsov^{1,2}

¹ *International Center for Theoretical Physics, P.O. Box 586, 34100 Trieste, Italy*

² *Landau Institute for Theoretical Physics, Kosygina str. 2, 117940 Moscow, Russia*

³ *Institute for Low Temp. Physics and Engineering, Lenin ave. 44, 310164 Kharkov, Ukraine.*

(February 1, 2008)

The level curvature distribution function is studied beyond the random matrix theory for the case of T-breaking perturbations over the orthogonal ensemble. The leading correction to the shape of the level curvature distribution is calculated using the nonlinear sigma-model. The sign of the correction depends on the presence or absence of the global gauge invariance and is different for perturbations caused by the constant vector-potential and by the random magnetic field. Scaling arguments are discussed that indicate on the qualitative difference in the level statistics in the dirty metal phase for space dimensionalities $d < 4$ and $d > 4$.

PACS numbers: 71.25.-s, 72.15.Rn, 05.45.+b

Since the seminal work by Thouless [1] the sensitivity of spectrum $\{E_n\}$ of disordered conductors to a small twist of the boundary conditions $\Psi(\rho, \mathbf{z} = \mathbf{0}) = e^{i\varphi} \Psi(\rho, \mathbf{z} = \mathbf{L})$ is considered as a powerful tool to probe the space structure of eigenfunctions and distinguish between the extended and the localized states. More precisely, the quantity K_n which is now referred to as the "level curvature", was introduced in Ref.[1] in order to describe this sensitivity quantitatively:

$$K_n = \Delta^{-1} \left. \frac{\partial^2 E_n(\varphi)}{\partial \varphi^2} \right|_{\varphi=0}, \quad (1)$$

where the mean level spacing $\Delta = (\nu L^d)^{-1}$ is related with the mean density of states ν and the size of the d -dimensional sample L .

In complex or disordered quantum systems, the quantity K_n fluctuates over the ensemble of energy levels $\{E_n\}$ or, for a given level, over the ensemble of realizations of disorder. The typical width of the distribution of level curvatures $P(K)$ is of the order of the dimensionless conductance $g = D/(L^2 \Delta)$, where D is the diffusion coefficient [1]. Thus studying this distribution one can obtain an information about the transition from metal to insulator with increasing the disorder (Anderson transition).

In the recent works [2] it has been shown that the distribution of the fluctuating quantity K_n is a particular example of *parametric level statistics*, i.e. statistics of spectral responses of the system to any perturbations proportional to some parameter φ .

A remarkable property of the parametric level statistics [2] similar to that of the usual level statistics [3,4], is that in a certain limit they are universal for all clas-

sically chaotic and disordered systems and can be described by the random matrix theory (RMT) of Wigner and Dyson [3,4]. For disordered systems considered here this limit coincides [5] with $g \rightarrow \infty$. For chaotic systems the same role is played [6] by the ratio $g = \gamma_1/\Delta$, where γ_1 is the first non-zero mode in the spectrum of the Perron-Frobenius operator that describes the chaotic behavior of the corresponding classical system. In particular, for the time-reversal-invariant system without spin-dependent interactions (orthogonal ensemble) the distribution of level curvatures, Eq.(1), was found [7,8] in this limit to have the form:

$$P_{WD}(k) = \frac{1}{2(1+k^2)^{\frac{3}{2}}}, \quad k = \frac{K}{\langle |K| \rangle}, \quad (2)$$

where $\langle |K| \rangle_{RMT} = 2g$ is the average modulus of the level curvature. Further study [9] showed that the form, Eq.(2), remains the same if weak localization is taken into account, only the dimensionless conductance in the expression for $\langle |K| \rangle$ is changed appropriately.

The form, Eq.(2), is universal. It does not depend, e.g. on details of the system and the perturbation. The only important point is that the system is T-invariant with $T^2 = 1$ and the perturbation breaks this invariance. However, the T-breaking perturbations can be of two different types. One of them (we call such perturbations "global") is represented by the twist of the boundary conditions or by the magnetic flux in the problem of persistent current [10]. In this case wave functions should have a significant amplitude on both boundaries $z = 0$ and $z = L$ in order the energy spectrum E_n be sensitive to the phase φ . Apparently, it is not so for the localized states, for which the distribution function $P(k)$ shrinks dramatically and is approximately of the logarithmically-normal form [11-13].

Another type of T-breaking perturbations is represented by the magnetic impurities or random magnetic fluxes. We call them "local perturbations", since they probe wave functions locally. Even for a localized state there should be a significant spectral response to changing the strength φ of such a local perturbation. So, one would expect neither any dramatic contraction nor the logarithmically-normal form of the distribution function $P(K)$ in this case.

Thus we conclude that there is no universality of the distribution function $P(k)$ in the Anderson insulator. It is reasonable to assume that the universality is broken

already at small but finite values of the parameter $1/g$ when the wave functions are still extended but have a pronounced structure. In the presence of such a structure, the system cannot be described any more by the basis-invariant ensemble of random matrices which is the main assumption of the random-matrix theory of Wigner and Dyson [3,4]. In order to use the curvature distribution function as a tool to probe the space structure of wave functions one should go beyond the random matrix theory.

In the present Letter we address the problem of corrections to the *shape* of the distribution function $P(k)$. A general approach to calculate such corrections using the nonlinear supersymmetric sigma-model [5] has been suggested in [14] and applied to distributions of different quantities [9,14,15]. It is based on the perturbative consideration of the non-zero diffusion modes which are integrated out to produce corrections to the zero-mode supersymmetric sigma-model [5]. The latter is then integrated exactly using the Efetov's parametrization [5].

Before going into details of the calculations we would like to formulate the main result and its consequences. The main result is the correction to the level curvature distribution function $P(k)$ for the case of T-breaking perturbations over the orthogonal ensemble:

$$\delta P(k) = \frac{C(d)}{(\pi g)^2} \frac{2 - 11k^2 + 2k^4}{2(1 + k^2)^{7/2}}, \quad k = \frac{K}{\langle |K| \rangle} \ll g, \quad (3)$$

where

$$C(d) = \sum_{\mathbf{q} \neq 0} \frac{1}{(\mathbf{q}^2)^2} \times \begin{cases} \left(\frac{4}{d} - 1\right), & \text{case I} \\ -1, & \text{case II} \end{cases} \quad (4)$$

Here $\mathbf{q} = \{q_1, \dots, q_d\}$, where $q_i = 2\pi n_i$, ($n_i = 0, \pm 1, \pm 2, \dots$) in the case of the periodic boundary conditions (for an unperturbed system) considered in this Letter.

A remarkable feature of Eqs.(3)-(4) for $d < 4$ is that the *sign* of the correction is different for global (case I) and local (case II) T-breaking perturbations. This reflects the principal difference between them which we discussed above. The positive sign of the correction at small K for the case of the global perturbations has been observed in the recent numerical simulations [12] on the 3D Anderson model. It implies the tendency towards the formation of a sharp peak in $P(K)$ at $K = 0$ which should be present in the insulator for the case of global perturbations and absent in the case of local ones.

Next we note that the sum in Eq.(4) is convergent only for $d < 4$. For $d > 4$ there is a problem of the upper momentum cut-off that cannot be done accurately within the framework of the nonlinear sigma-model approach. It turns out that for global perturbations the coefficient $C(d)$ changes sign precisely at $d = 4$ where the difficulty with the divergent sum is first encountered. This coincidence could hardly be accidental. It may indicate on the qualitative difference in the structure of wave functions in disordered systems for $d < 4$ and $d > 4$, especially near

the Anderson transition where the parameter $1/g^2$ is not very small. Meanwhile the problem of the Anderson transition in higher dimensions is becoming a physical issue because of some models [16] that map the problem of disordered systems of *interacted* electrons onto the Anderson model in higher dimensions.

Now we outline the derivation of the announced results. The first step is to express the level curvature distribution function $P(K)$ in terms of the two-level correlation function $R(\omega, \varphi)$. This is done in the same way as in Ref.[17]:

$$P(K) = \lim_{\varphi \rightarrow 0} \left[\frac{\varphi^2}{2} R \left(\omega = \frac{\varphi^2}{2} K \Delta; \varphi \right) \right]. \quad (5)$$

The second step is to represent the two-level correlation function in Eq.(5) in terms of the supersymmetric nonlinear sigma-model which has been derived in Ref.[5] from the model of non-interacting electrons in a random potential. Using the results of Ref.[5] we arrive at:

$$P(K) = - \frac{\varphi^2}{16} \Re \frac{\partial^2}{\partial j_+ \partial j_-} \int \mathcal{D}Q e^{-F_\varphi[Q; K, j_\pm]} \Big|_{\varphi, j_\pm \rightarrow 0}. \quad (6)$$

In this equation $Q(\mathbf{r})$ is the 8×8 supermatrix field of a certain symmetry which obeys the constraint $Q^2 = 1$, and $F_\varphi[Q; K, j_\pm] = F[Q; K] + F[Q; j_+] + F[Q; j_-]$, where:

$$F[Q; j_\pm] = j_\pm \int \text{Str}[P_\pm P_B Q] \frac{d\mathbf{r}}{L^d} \quad (7)$$

Here $\text{Str}[A] = \text{tr}[A_{FF}] - \text{tr}[A_{BB}]$ stands for the supertrace, $P_\pm$ are the projectors onto the retarded (RR) or advanced (AA) sectors and P_B is the projector onto the boson-boson sector of the superfield Q (see Ref.[5] for details and definitions). An equivalent representation has been used in Ref.[9].

The main information about the underlying physics is contained in the action $F[Q; K]$. For global perturbations it takes the form:

$$F_\varphi[Q; K] = \frac{\pi}{8} g L^2 \int \text{Str} \left(\vec{\nabla} Q - \frac{i\vec{\varphi}}{L} [Q, \tau] \right)^2 \frac{d\mathbf{r}}{L^d} + i \frac{\pi}{8} K \varphi^2 \int \text{Str}[\Lambda Q] \frac{d\mathbf{r}}{L^d}, \quad (8)$$

where $\vec{\varphi} = (0, \varphi)$ is supposed to be directed along the z -axis; $\Lambda = P_+ - P_-$, and $\tau = \sigma_z \otimes P_+$ is proportional to the Pauli matrix σ_z that breaks the symmetry in the quaternionic subspace of the superfield Q and reflects the T-symmetry breaking.

For local T-breaking perturbations the linear in φ term in Eq.(8) is absent.

The representation, Eqs.(6)-(8), in terms of the field $Q(\mathbf{r})$ contains all the spatial diffusion modes $\gamma_q = (D/L^2)\mathbf{q}^2$. However, in doing the limit $\varphi \rightarrow 0$ in Eq.(6) the main

role is played by the zero mode which corresponds to $\mathbf{q} = 0$. At $\varphi = 0$ this mode does not cost any energy no matter how large are the components of the field Q in the non-compact boson-boson sector [5]. It is the arbitrary large amplitudes of the zero mode components of the field Q that compensate the infinitesimal parameter φ in Eqs.(6)-(8) and lead to a finite result for $P(K)$. Thus the space independent zero mode Q_0 must be considered non-perturbatively. In the limit $g \rightarrow \infty$ all the non-zero modes can be neglected [5], and one arrives [9] at the RMT result, Eq.(2). For finite $1/g$ the non-zero modes should be also taken into account. However, all the non-zero modes can be treated perturbatively for $g \gg 1$ to lead to some corrections to the zero-mode action. In order to obtain these corrections we have to separate zero modes from all other modes and then integrate over all the non-zero modes using a certain perturbative scheme. This is done by means of the transformation [14]:

$$Q(\mathbf{r}) = V_0^{-1} \tilde{Q}(\mathbf{r}) V_0, \quad Q_0 = V_0^{-1} \Lambda V_0. \quad (9)$$

The field $\tilde{Q}(\mathbf{r})$ that must also obey the constraint $\tilde{Q}^2 = 1$, is parametrized as follows [14]:

$$\tilde{Q} = (1 - W/2)\Lambda(1 - W/2)^{-1}, \quad (10)$$

where the supermatrix $W(\mathbf{r})$ does not contain the zero spatial mode.

The transformation from $Q(\mathbf{r})$ to new variables $Q_0, W_{\mathbf{q} \neq 0}$ involves [15] a non-trivial Jacobian $J = e^{-F_J[W]}$. To the first non-vanishing order in W it is given by [18]:

$$F_J[W] = -\frac{1}{8} \int Str[W^2] \frac{d\mathbf{r}}{L^d}. \quad (11)$$

In order to integrate over W perturbatively, one has to expand the action, Eqs.(7),(8),(11) in powers of W separating the V_0 -independent quadratic in W part $F_0[W]$ and consider all the rest W -dependent terms as perturbations using the Wick's theorem.

As a result of such calculations we have obtained the correction up to the $1/g^2$ order to the zero-mode action $F[Q_0; K, j_{\pm}]$. It is too lengthy to present it here and contains all possible linear and bi-linear combinations of vertices encountered in $F[Q_0; K, j_{\pm}]$ plus 4 new vertices.

A remarkable feature of the Efetov's parametrization [5] that has not been mentioned so far is that in the limit of large "non-compact angles" $\lambda_{1,2} \rightarrow \infty$, the new vertices reduce to the product of the "old" ones. Thus up to the leading order of $\lambda_{1,2}^4$ we found:

$$\begin{aligned} Str[(\Lambda Q_0)^2] &= -[Str[\Lambda Q_0]]^2, \\ Str[(\tau Q_0)^4] &= -\frac{1}{2} [Str[(\tau Q_0)^2]]^2 \\ Str[(\tau Q_0)^2 \Lambda Q_0] &= -\frac{1}{2} Str[(\tau Q_0)^2] Str[\Lambda Q_0], \\ Str[(\tau Q_0)^2 P_{\pm} P_B Q_0] &= -\frac{1}{2} Str[(\tau Q_0)^2] Str[P_{\pm} P_B Q_0]. \end{aligned} \quad (12)$$

This circumstance simplifies the solution dramatically, since it allows to obtain all the corrections to $P(K)$ just by differentiating of the RMT result, Eq.(2), over g and K . Since the K and g -dependences of the RMT distribution function $\tilde{P}_{WD}(K) = (2g)^{-1} P_{WD}(K/2g)$ are closely related, the result can be formulated in terms of the g -derivatives alone. It is convenient to introduce the parameter $\alpha = \ln g$. Then the corrections to the distribution function $P(K)$ can be represented in the form:

$$\begin{aligned} \delta P(K) &= -\frac{1}{\pi g} \sum_{\mathbf{q} \neq 0} \frac{1}{\mathbf{q}^2} \left(\frac{\partial \tilde{P}_{WD}(K)}{\partial \alpha} \right) + \\ &+ \frac{1}{(\pi g)^2} \sum_{\mathbf{q} \neq 0} \frac{1}{(\mathbf{q}^2)^2} \left[\frac{9}{2} - \frac{16}{d} + \frac{36}{d(d+2)} \right] \frac{\partial \tilde{P}_{WD}(K)}{\partial \alpha} + \\ &+ \frac{1}{(\pi g)^2} \sum_{\mathbf{q} \neq 0} \frac{1}{(\mathbf{q}^2)^2} \left(\frac{4}{d} - 1 \right) \frac{\partial^2 \tilde{P}_{WD}(K)}{\partial \alpha^2}. \end{aligned} \quad (13)$$

The result, Eq.(13), holds for the global perturbations. For such perturbations ∇Q enters Eq.(8) as a "covariant derivative" $\nabla Q - \frac{i\varphi}{2} [Q, \tau]$. As a result, the linear in φ "cross"-vertex $\vec{\varphi} Str[Q \vec{\nabla} Q \tau]$ appears in the action. It is this vertex that leads to terms in Eq.(13) proportional to $1/d$ or $1/d(d+2)$. For local perturbations such terms are absent, since the linear in φ vertex does not appear at all [5].

The term with the second derivative in Eq.(13) contributes to the correction to the *shape* of the distribution function $P(k)$ and leads to the announced result, Eq.(3).

The terms proportional to the first derivative can be absorbed into the RMT distribution function $P_{WD}(k)$ by changing the average $\langle |K| \rangle$. The $1/g$ correction to this value first found in Ref.[9] is nothing but the usual weak-localization correction to the quantity $\langle |K| \rangle_{RMT} = 2g$. Thus up to the first order in the inverse "bare" conductance $1/g$, the quantity $\frac{1}{2} \langle |K| \rangle$ behaves exactly like the true dimensionless conductance $\bar{g}$.

Moreover, it turns out that all the $1/g^2$ corrections to $P(K)$ proportional to $(\sum \frac{1}{\mathbf{q}^2})^2$ cancel each other in the same way as in the dimensionless conductance $\bar{g}$ [20]. This is in a full agreement with the one-parameter scaling [21] extended to describe the distributions of mesoscopic fluctuations [19]. In the region where the one-parameter scaling works, the distribution functions of all quantities (measured in proper units) should depend only on one additional parameter - the average dimensionless conductance $\bar{g}$ that absorbs all microscopic parameters of the system and includes all quantum corrections proportional to the powers of $g^{-1} \sum \frac{1}{\mathbf{q}^2}$. For the case of the level curvature distribution it means that $P(K) = P(K; \bar{g})$, where $\bar{g}$ should obey the scaling equation [21] that does not allow [20] the term proportional to $g^{-2} (\sum \frac{1}{\mathbf{q}^2})^2$ in the perturbative expansion of $\bar{g}/g$ in powers of $1/g$.

From this viewpoint the corrections to $P(K)$ proportional to $g^{-2} \sum \frac{1}{(\mathbf{q}^2)^2}$ should be considered as the *main*

terms in the expansion of $P(K; \bar{g})$ in powers of the *true* inverse dimensionless conductance $1/\bar{g}$ renormalized by the localization effects rather than the second-order terms of an expansion in powers of the "bare" value of $1/g$. With due regard of these corrections the shape of the distribution function $P(k)$ deviates from the RMT result and the average $\langle |K| \rangle$ is no more proportional to the average dimensionless conductance:

$$\langle |K| \rangle = 2\bar{g} \left[1 + \frac{A(d)}{(\pi\bar{g})^2} \sum_{\mathbf{q} \neq 0} \frac{1}{(\mathbf{q}^2)^2} \right], \quad (14)$$

where $A(d) = 9/2$ for the local perturbations and $A(d) = (9/2 - 16/d + 36/d(d+2))$ for the global perturbations.

In conclusion we would like to discuss a consequence of the form of corrections to different level statistics for the applicability of the RMT results. The typical relative correction [see e.g. [14,22]] has the same form as Eq.(14) and is proportional to $\bar{g}^{-2} \sum \frac{1}{(\mathbf{q}^2)^2}$. Let us suppose that the system is close enough to the Anderson transition (from the metal side) but it is still not in the critical region, so that $\pi\bar{g} \gg 1$ but it is not very large. Then for $d < 4$, where the sum in Eq.(14) is convergent, the correction is still small, and the RMT can be considered as a good zero approximation. However, it is not the case for $d > 4$ if $\pi\bar{g} < (L/r_0)^{d/2-2}$, where r_0^{-1} is the upper momentum cut-off. In the scaling picture of the Anderson transition [21] the dimensionless conductance $\bar{g}$ can be made arbitrary close to its critical value $g_c \sim 1$ by the fine tuning of the elastic scattering mean free path $l \approx l_c$ at a constant $L \gg l \sim r_0$. Thus we conclude that the above condition is always possible to achieve for $d > 4$ at some strength of disorder $W_{c1}(L)$ which is less than the critical one W_c . For $W > W_{c1}$ the RMT results do not work even as a zero approximation for the level statistics.

In summary, we have calculated the leading corrections to the shape of the level curvature distribution function $P(K)$ beyond the random matrix theory. For T-breaking perturbations over the orthogonal ensemble, the corrections are not universal and depend qualitatively on the presence or absence of the global gauge invariance. In both cases the average $\frac{1}{2}\langle |K| \rangle$ shows deviations from the average quantum dimensionless conductance $\bar{g}$. The corrections are in agreement with the one-parameter scaling. The consequence of the scaling hypothesis and the form of the corrections is a qualitative difference in level statistics for space dimensionalities above and below $d = 4$. Below $d = 4$ the RMT is a good zero approximation for level statistics in the metal phase, while for $d \geq 4$ there should exist a critical value of disorder $W_{c1} < W_c$ above which RMT fails to describe spectral correlations.

Acknowledgements We thank B.L.Altshuler, Chaitali Basu, C.M.Canali, V.I.Fal'ko, Y.V.Fyodorov, I.V.Lerner and A.D.Mirlin for stimulated discussions. Support from RFBR and INTAS grant RFBR/INTAS 95-675 (V.E.K.) is gratefully acknowledged.

-
- [1] D. J. Thouless, Phys. Rep. **13**, 93 (1974).
 - [2] B. D. Simons and B. L. Altshuler, Phys. Rev. B. **48**, 5422 (1993); A. Szafer and B. L. Altshuler, Phys. Rev. Lett. **70**, 587 (1993); B. D. Simons, P. A. Lee and B. L. Altshuler, Phys. Rev. Lett. **70**, 4122 (1993).
 - [3] E. P. Wigner, Proc. Cambridge Philos. Soc. **47**, 790 (1951); F. J. Dyson, J. Math. Phys. **3**, 140 (1962).
 - [4] M. L. Mehta, *Random matrices* (Academic Press, Boston, 1991).
 - [5] K. B. Efetov, Adv.Phys. **32**, 53 (1983).
 - [6] A. V. Andreev, O. Agam, B. D. Simons, and B. L. Altshuler, Phys. Rev. Lett. **76**, 3947 (1996).
 - [7] J. Zakrewski and D. Delande, Phys. Rev. E. **47**, 1650 (1993).
 - [8] F. von Oppen, Phys. Rev. Lett. **73**, 798 (1994).
 - [9] Y. V. Fyodorov and H.- J. Sommers, Phys. Rev. E. **51**, R2719 (1995).
 - [10] M. Buttiker, Y. Imry, and R. Landauer, Phys. Lett. A. **96**, 365 (1983).
 - [11] K. Zyczkowski, L. Molinari, and F. M. Izrailev, J. Phys. I France. **4**, 1469 (1994).
 - [12] C. M. Canali, C. Basu, W. Stephan and V. E. Kravtsov, Phys. Rev. B. **54**, 1431 (1996).
 - [13] M. Titov, and Y. V. Fyodorov, (unpublished).
 - [14] V. E. Kravtsov and A. D. Mirlin, Pis'ma Zh. Exp. Teor. Fiz. **60**, 645 (1994) [JETP Lett. **60**, 656 (1994)].
 - [15] Y. V.Fyodorov and A. D. Mirlin, Phys. Rev. B **51**, 13403 (1995).
 - [16] B. L. Altshuler, Y. Gefen, A. Kamenev and L. Levitov (unpublished).
 - [17] V. E. Kravtsov and M. R. Zirnbauer, Phys. Rev. B. **46**, 4332 (1992).
 - [18] There was a minor mistake in calculating the Jacobian in Ref.[15]. The correct answer is given by Eq.(11)
 - [19] B. L. Altshuler, V. E. Kravtsov, and I. V. Lerner, Zh. Eksp. Teor. Fiz. **91**, 2276 (1986) [Sov. Phys. JETP **64**, 1352 (1986)].
 - [20] L. P. Gor'kov, A. I. Larkin, and D. E. Khmelnitskii, Pis'ma v ZhETF **30**, 248 (1979) [Sov. Phys. JETP Letters **30**, 228 (1979)].
 - [21] E. Abrahams, P. W. Anderson, D. C Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
 - [22] A. V. Andreev and B. L. Altshuler, Phys. Rev. Lett. **75**, 902 (1995).